

# GEOMETRIZED VACUUM PHYSICS. PART I: ALGEBRA OF STIGNATURES

## FÍSICA DEL VACÍO GEOMETRIZADA. PARTE I: ALGEBRA DE STIGNATURES

---

**Mikhail Batanov-Gaukhman<sup>1</sup>**

(1) Moscow Aviation Institute (National Research University), Institute No. 2 "Aircraft and rocket engines and power plants", st. Volokolamsk highway, 4, Moscow – Russia, 125993  
(e-mail: alsignat@yandex.ru)

*Recibido: 02/11/2022 - Evaluado: 04/12/2022 - Aceptado: 26/12/2022*

---

### ABSTRACT

This article is the first part of a scientific project under the general title "Geometrized Vacuum Physics". This study is based on two main postulates: 1) independence of the propagation velocity of electromagnetic waves in vacuum from their frequency; 2) the constancy of the averaged zero vacuum balance, associated with the assertion that only mutually opposite formations are born from vacuum, so that on average they completely compensate for each other's manifestations. In this part of the "Geometrized Vacuum Physics" the foundations of the Algebra of Stignatures are laid, which is the mathematical and logical foundation of the entire project. In the next articles of this project it will be shown that the Algebra of Stignatures can be used for the development of "zero" (vacuum) technologies, algebraic genetics, vacuum cosmology, the vacuum standard model of "elementary particles", vacuum gravity and levitation, vacuum energy, ethics and aesthetics and many other branches of knowledge. At the end of this article, one of the many possible applications of the Algebra of Stignatures is given, in particular, the basics of the signature-spectral analysis are outlined, with the help of which the possibilities of communication channels can be significantly expanded.

### RESUMEN

Este artículo es la primera parte de un proyecto científico bajo el título general "Física del vacío geometrizada". Este estudio se basa en dos postulados principales: 1) independencia de la velocidad de propagación de las ondas electromagnéticas en el vacío de su frecuencia; 2) la constancia del balance de vacío promediado cero, asociado con la afirmación de que solo las formaciones mutuamente opuestas nacen del vacío, de modo que en promedio compensan completamente las manifestaciones de cada uno. En esta parte de la "Física Geometrizada del Vacío" se sientan las bases del Álgebra de Stignatures, que es el fundamento matemático y lógico de todo el proyecto. En los próximos artículos de este proyecto se mostrará que el Álgebra de Stignatures puede ser utilizada para el desarrollo de tecnologías de "cero" (vacío), genética algebraica, cosmología de vacío, el modelo estándar de vacío de "partículas elementales", gravedad de vacío y levitación, energía del vacío, ética y estética y muchas otras ramas del saber. Al final de este artículo, se proporciona una de las muchas aplicaciones posibles del Stignature Algebra, en particular, se describen los conceptos básicos del análisis espectral de firmas, con la ayuda de la cual las posibilidades de los canales de comunicación pueden ampliarse significativamente.

Keywords: vacuum, algebra of stignatures, vacuum balance, signature  
Palabras clave: vacío, álgebra de stignatures, balance de vacío, signatura

## INTRODUCTION

In modern physics, there are (Shipov, 1998; Chambers, 2004; Roberts., 2000; Barrow, 2002; Di Teodoro & Contreras, 2023): technical vacuum (discharged gas); physical vacuum (the lowest energy state of a set of scalar, vector, tensor and spinor quantum fields); Einstein vacuum (generally a curved 4-dimensional space-time continuum surrounding neutral or charged physical bodies); ideal vacuum (emptiness, i.e. 3-dimensional space, in which any curvature and particles are completely absent).

This first part of "Geometrized Vacuum Physics" focuses on the ideal vacuum in order to create a mathematical apparatus "Algebra of Stignatures", suitable for the study of vacuum phenomena and the development of "zero" (vacuum) technologies. In subsequent articles of this project, a description of curved sections of vacuum and stable vacuum formations is expected. The concept of "signature" is introduced in paragraph 5.17 of this article. The fundamentals of the Algebra of Signatures and the explanation of this term will be presented in the next article of the proposed project.

First, we abstract from all kinds of vacuum processes, and consider the local region of a completely flat empty 3-dimensional space (i.e., ideal vacuum).

This work is based on three reliable (experimentally confirmed) facts:

- 1) all known electromagnetic waves, regardless of their oscillation frequency, propagate in vacuum with the same speed of light  $c = 299,792,458$  m/s (fixed by the resolution of the 26-th General Conference on Weights and Measures, based on CODATA data). Despite the possible instability of the properties of emptiness, the postulate of the constancy of the speed of light in vacuum is accepted in this work as an initial axiom. At the same time, as will become clear below, the mathematical apparatus of the Algebra of Stignatures turned out to be independent of the propagation velocity of perturbations. Therefore, the Algebra of Stignatures is universal and applicable to the study of any 3-dimensional medium;
- 2) all averaged characteristics of an average flat area of vacuum (momentum, angular momentum, spin, etc.) are equal to zero;
- 3) if something is born from vacuum, then it must be in a mutually opposite form (particle - antiparticle, convexity - concavity, wave - antiwave, etc.). This property of vacuum is referred to in this work as "averaged zero vacuum balance" or "vacuum balance" for short.

The foundations of the Algebra of Stignatures developed in this work are proposed as a universal mathematical apparatus suitable for studying not only the properties of vacuum, but also any other liquid, solid and gaseous continuous media in which wave disturbances propagate at a constant speed. Also, the Algebra of Stignatures can be useful for philosophical rethinking of the knowledge of ancient civilizations and for application in many branches of scientific knowledge, for example, in: coding theory, algebraic genetics, vacuum energy, elementary particle physics, etc.

## FUNDAMENTALS AND METHODOLOGY

### *1 Longitudinal stratification of ideal vacuum into $\lambda_{m,n}$ -vacuums*

Consider a 3-dimensional volume of an ideal vacuum, in which there are no particles and no curvature. In what follows, for brevity, the ideal vacuum will be referred to as "vacuum".

Let's probe this volume of "vacuum" with laser beams from three mutually perpendicular directions, so that the beams form a 3-dimensional cubic lattice (Fig. 1*a,b*).

Rays of light in a vacuum are not visible, but they can be visualized with a finely dispersed sol. Of course, a "vacuum" filled with a sol is not a perfect vacuum. Nevertheless, the rays propagate in the "vacuum" itself (i.e.,

between the particles of the sol), while the influence of the sol on the metric-dynamic properties of the macroscopic volume of the "vacuum" in the case under consideration can be neglected. In addition, if the sol is removed from this "vacuum" volume, then the rays will still remain in it, although they will not be visible.




Fig. 1: a) Laser beams of light in vacuum, visualized with a fine sol; b) A 3-dimensional lattice in a "vacuum" consisting of mutually perpendicular monochromatic light rays with a wavelength  $\lambda_{m,n}$  while the edge length of a given cubic cell is  $\varepsilon_{m,n} \sim 10^2 \lambda_{m,n}$

Let, for example, to probe of "vacuum" are used laser beams of light (i.e. narrowly directed monochromatic electromagnetic oscillations) with a wavelength  $\lambda_{4,-5}$ , taken from the range of lengths  $\Delta\lambda = 10^{-4} \div 10^{-5}$  cm. Then we get a 3-dimensional lattice consisting of such mutually perpendicular laser beams, with the edge length of one cubic cell  $\varepsilon_{4,-5} \sim 100 \cdot \lambda_{4,-5}$  (see Fig.1b). This cubic lattice will be called  $\lambda_{4,-5}$ -vacuum (or 3D<sub>4,-5</sub>-landscape).

We divide the entire range of electromagnetic (light) wave lengths into a set of subranges  $\Delta\lambda = 10^m \div 10^n$  cm, where  $n = m + 1$  ( $m$  and  $n$  are integers). Then, in the same way as shown in Fig. 1, we will probe the studied volume of the "vacuum" with monochromatic light rays with wavelengths  $\lambda_{m,n}$  from all subranges  $\Delta\lambda = 10^m \div 10^n$  cm. As a result, we obtain an almost infinite number of nested  $\lambda_{m,n}$ -vacuums (i.e. 3D<sub>m,n</sub>-landscapes) with the corresponding lengths of edges of cubic cells  $\varepsilon_{m,n} \sim 100 \lambda_{m,n}$  (see Fig. 2).




Fig. 2: Discrete set of nested  $\lambda_{m,n}$ -vacuums of the same 3-dimensional void volume, where  $\lambda_{m,n} > \lambda_{m+1,n+1} > \lambda_{m+2,n+2} > \lambda_{m+3,n+3} \dots$

The value of the edge of the cubic cell of each  $\lambda_{m,n}$ -vacuum

$$\varepsilon_{m,n} \sim 10^2 \lambda_{m,n} \tag{1}$$

follows from the condition of applicability of geometric optics  $\lambda_{m,n} \rightarrow 0$ , i.e. when the thickness of the light beam is much smaller than the value of the corresponding cubic cell, while the beam thickness can be neglected.

The question remains open: – “Are there any restrictions on the frequency  $\omega$  or the wavelength  $\lambda$  of the electromagnetic wave, both in the direction of their increase, and in the direction of decrease? If the critical values:  $\omega_{max} = 2\pi c/\lambda_{max}$  and  $\omega_{min} = 2\pi c/\lambda_{min}$  exist, these will be very important characteristics of the vacuum. Today, as far as the author knows, the frequency range of observed electromagnetic waves extends from 2 Hz to  $10^{20}$  Hz, while no restrictions on the expansion of this range have been experimentally found.

### 2 Geodesic lines of the curved section $\lambda_{m,n}$ -vacuum

Long-term experimental data show that monochromatic light rays in the entire observed wavelength range  $\Delta\lambda$  propagate in "vacuum" at the same speed of light  $c$  and according to the same laws of electrodynamics. Therefore, if the region of vacuum under study is not curved, then all  $\lambda_{m,n}$ -vacuums (i.e.,  $3D_{m,n}$ -landscapes) will be represented as ideal cubic lattices (see Figs. 1 and 2), because the geodesic lines of all these non-curved  $\lambda_{m,n}$ -vacuums are direct rays of light. In this case,  $\lambda_{m,n}$ -vacuums will differ from each other only by the length of the edge of the cubic cell  $\varepsilon_{m,n} \sim 10^2\lambda_{m,n}$  (see Fig. 2).

However, if the region of vacuum under study turns out to be curved, then all  $\lambda_{m,n}$ -vacuums will differ somewhat from each other due to the fact that light rays with different wavelengths have different thicknesses. This circumstance is theoretically substantiated in the sections of geometric optics related to the resolution of optical devices (Tyurin *et al.*, 2009; Peatross & Ware, 2015) and is confirmed by experimental data (see Fig. 3).




Fig. 3: Experimental data on the thickness of the laser beam depending on the length wave  $\lambda$  of the corresponding monochromatic radiation

In this case, each  $\lambda_{m,n}$ -vacuum (i.e.,  $3D_{m,n}$ -landscape) will be unique (see Fig. 4), since vacuum irregularities are averaged within the thickness of the probing light beam.




Fig. 4: Illustration of a curved  $\lambda_{m,n}$ -vacuum embedded in curved  $\lambda_{f,d}$ -vacuum (where  $\lambda_{f,d} > \lambda_{m,n}$ )

Therefore, one  $\lambda_{m,n}$ -vacuum is only one 3-dimensional "slice" of the curved vacuum region. For a more complete description of the curved section of the vacuum, it is necessary to have an infinite set of curved  $\lambda_{m,n}$ -vacuums nested in each other.

Thus, the local volume of vacuum is an infinitely complex system consisting of an infinite number of nested  $\lambda_{m,n}$ -vacuums. However, the situation is simplified by the fact that in the entire studied range of electromagnetic wave lengths, all  $\lambda_{m,n}$ -vacuums obey the same physical laws. Therefore, the knowledge gained in the study of one  $\lambda_{k,r}$ -vacuum is automatically extended to all other  $\lambda_{m,n}$ -vacuums.

Below, the mathematical apparatus of the Algebra of Stignatures is developed, designed to study the local volume of only one  $\lambda_{m,n}$ -vacuum. But this apparatus is suitable for studying not only all  $\lambda_{m,n}$ -vacuums, but also any other continuous media in which wave disturbances propagate at a constant speed.

### 3 Sixteen rotating 4-bases

Let's return to the consideration of the undistorted volume of one of the  $\lambda_{m,n}$ -vacuums (see Fig. 1) and examine the "vacuum" region in the vicinity of the point  $O$  (see Fig. 5).




Fig. 5: Non-curved 3D light lattice of  $\lambda_{m,n}$ -vacuum, revealed from the "vacuum" (emptiness) by means of mutually perpendicular monochromatic rays of light with a wavelength  $\lambda_{m,n}$ . The cells of such a lattice are cubes with edge length  $\epsilon_{m+n} \sim 10^2 \lambda_{m,n}$

We calculate how many orthogonal 3-bases originate at the central point  $O$  (see Fig. 5). If we spread the 3-bases from the point  $O$  in different directions, then it turns out that there are 16 of them (Fig. 6 *a,b*).




Fig. 6: Sixteen 3-bases at the central point  $O$  of the studied volume of  $\lambda_{m,n}$ -vacuum. *a)* 8 internal 3-bases; *b)* 8 outer 3-antibases; *c)* adjacent 3-bases

Eight 3-bases refer to the cubic cell itself (Fig. 6*a*), and eight opposite 3-antibases refer to adjacent cubic cells (Fig. 6*b,c*).

According to the "vacuum balance" condition, any movement in a vacuum must be accompanied by a similar anti-motion. Therefore, if one 3-basis (together with the cubic cell) is rotated clockwise, then this is only possible if the adjacent cubic cell (together with the 3-antibasis) is similarly rotated counterclockwise, since there is no fulcrum in vacuum.

In connection with the above, it is convenient to add to all eight 3-bases (Fig. 6a) along the fourth time axis  $t$ , and to eight 3-antibases (Fig. 6b) add along the fourth anti-axis (i.e., oppositely directed axis) of time  $-t$ .

The time axis  $t$  is determined by the angular frequency of rotation of the 3-basis (ie, the number of revolutions per unit time). The rotation of the 3-basis with a constant angular velocity is described by the expression  $d\varphi/dt = \omega$  (where  $\varphi$  and  $\omega$  are the phase and angular frequency of rotation of the 3-basis). Integrating this expression, we get the time axis  $t = \varphi / \omega$ . The rotation of the 3-antibasis in the opposite direction similarly forms the anti-axis of time  $-t = \varphi / \omega$ .

Thus, at the considered point  $O$   $\lambda_{m,n}$ -vacuum (Fig. 5) there are  $8 + 8 = 16$  orthogonal 4-bases (Fig. 6), which are shown in Fig. 7.




Fig. 7: Sixteen 4-bases starting at point  $O$ , obtained by adding to eight 3-bases (Fig. 6a) of the fourth time axis and adding to eight 3-antibases (Fig. 6b) the fourth anti-time axis

#### 4 Stignature of an affine 4-dimensional space

Each of the sixteen 4-bases shown in Fig. 7 specifies the direction of the axes of a 4-dimensional affine (i.e., vectors) space. To introduce the characteristic "stignature" of an affine space, we first define the concept of "base".




Fig. 8: An affine 4-dimensional space, the directions of the axes of which are given by the 4-basis  $\mathbf{e}^5(\mathbf{e}_0^{(5)}, \mathbf{e}_1^{(5)}, \mathbf{e}_2^{(5)}, \mathbf{e}_3^{(5)})$  with the conventional signature  $\{+ + + +\}$

We choose from the sixteen 4-bases shown in Fig. 7, one 4-basis, for example,  $\mathbf{e}^5 (\mathbf{e}_0^{(5)}, \mathbf{e}_1^{(5)}, \mathbf{e}_2^{(5)}, \mathbf{e}_3^{(5)})$  and let's call it "base" (see Fig. 8).

We conditionally accept that the directions of all unit vectors of the "base" are positive (see Fig. 8)

$$\mathbf{e}^{(5)}(\mathbf{e}_0^{(5)}, \mathbf{e}_1^{(5)}, \mathbf{e}_2^{(5)}, \mathbf{e}_3^{(5)}) = (+1, +1, +1, +1) \rightarrow \{+ + + +\}. \tag{2}$$

Here we for the first time introduce the abbreviated notation  $\{+ + + +\}$ , which we will call the "stignature" of the affine space given by the 4-basis  $\mathbf{e}^5$  (that is, the "base"). That is, a stignature is a set of 4 signs that determine the direction of the axes of a 4-dimensional affine space in relation to the direction of the axes of the base affine space.

All other 4-bases shown in Fig. 7 have the following signatures with respect to the directions of the "base" unit vectors (i.e., the 4-basis  $\mathbf{e}^{(5)}$ ) (see Table 1).

Table 1: Stignatures of affine spaces

4-basis	Stignature	4-basis	Stignature
$\mathbf{e}^1 (\mathbf{e}_0^{(1)}, \mathbf{e}_1^{(1)}, \mathbf{e}_2^{(1)}, \mathbf{e}_3^{(1)}) =$ $= (1, 1, -1, 1) \rightarrow$	$\{+ + - +\}$	$\mathbf{e}^9 (\mathbf{e}_0^{(9)}, \mathbf{e}_1^{(9)}, \mathbf{e}_2^{(9)}, \mathbf{e}_3^{(9)}) =$ $= (-1, 1, -1, 1) \rightarrow$	$\{- + - +\}$
$\mathbf{e}^2 (\mathbf{e}_0^{(2)}, \mathbf{e}_1^{(2)}, \mathbf{e}_2^{(2)}, \mathbf{e}_3^{(2)}) =$ $= (1, -1, -1, -1) \rightarrow$	$\{+ - - -\}$	$\mathbf{e}^{10} (\mathbf{e}_0^{(10)}, \mathbf{e}_1^{(10)}, \mathbf{e}_2^{(10)}, \mathbf{e}_3^{(10)}) =$ $= (-1, 1, -1, -1) \rightarrow$	$\{- - - -\}$
$\mathbf{e}^3 (\mathbf{e}_0^{(3)}, \mathbf{e}_1^{(3)}, \mathbf{e}_2^{(3)}, \mathbf{e}_3^{(3)}) =$ $= (1, 1, -1, -1) \rightarrow$	$\{+ + - -\}$	$\mathbf{e}^{11} (\mathbf{e}_0^{(11)}, \mathbf{e}_1^{(11)}, \mathbf{e}_2^{(11)}, \mathbf{e}_3^{(11)}) =$ $= (-1, 1, -1, -1) \rightarrow$	$\{- + - -\}$
$\mathbf{e}^4 (\mathbf{e}_0^{(4)}, \mathbf{e}_1^{(4)}, \mathbf{e}_2^{(4)}, \mathbf{e}_3^{(4)}) =$ $= (1, -1, -1, 1) \rightarrow$	$\{+ - - +\}$	$\mathbf{e}^{12} (\mathbf{e}_0^{(12)}, \mathbf{e}_1^{(12)}, \mathbf{e}_2^{(12)}, \mathbf{e}_3^{(12)}) =$ $= (-1, -1, -1, 1) \rightarrow$	$\{- - - +\}$
$\mathbf{e}^{(5)} (\mathbf{e}_0^{(5)}, \mathbf{e}_1^{(5)}, \mathbf{e}_2^{(5)}, \mathbf{e}_3^{(5)}) =$ $= (1, 1, 1, 1) \rightarrow$	$\{+ + + +\}$	$\mathbf{e}^{13} (\mathbf{e}_0^{(13)}, \mathbf{e}_1^{(13)}, \mathbf{e}_2^{(13)}, \mathbf{e}_3^{(13)}) =$ $= (-1, 1, 1, 1) \rightarrow$	$\{- + + +\}$
$\mathbf{e}^6 (\mathbf{e}_0^{(6)}, \mathbf{e}_1^{(6)}, \mathbf{e}_2^{(6)}, \mathbf{e}_3^{(6)}) =$ $= (1, -1, 1, -1) \rightarrow$	$\{+ - + -\}$	$\mathbf{e}^{14} (\mathbf{e}_0^{(14)}, \mathbf{e}_1^{(14)}, \mathbf{e}_2^{(14)}, \mathbf{e}_3^{(14)}) =$ $= (-1, -1, 1, -1) \rightarrow$	$\{- - + -\}$
$\mathbf{e}^7 (\mathbf{e}_0^{(7)}, \mathbf{e}_1^{(7)}, \mathbf{e}_2^{(7)}, \mathbf{e}_3^{(7)}) =$ $= (1, 1, 1, -1) \rightarrow$	$\{+ + + -\}$	$\mathbf{e}^{(15)} (\mathbf{e}_0^{(15)}, \mathbf{e}_1^{(15)}, \mathbf{e}_2^{(15)}, \mathbf{e}_3^{(15)}) =$ $= (-1, 1, 1, -1) \rightarrow$	$\{- + + -\}$
$\mathbf{e}^8 (\mathbf{e}_0^{(8)}, \mathbf{e}_1^{(8)}, \mathbf{e}_2^{(8)}, \mathbf{e}_3^{(8)}) =$ $= (1, -1, 1, 1) \rightarrow$	$\{+ - + +\}$	$\mathbf{e}^{16} (\mathbf{e}_0^{(16)}, \mathbf{e}_1^{(16)}, \mathbf{e}_2^{(16)}, \mathbf{e}_3^{(16)}) =$ $= (-1, -1, 1, 1) \rightarrow$	$\{- - + +\}$

Stignatures given in Table. 1 are combined into a 16-component matrix:

$$stign(e_i^{(a)}) = \begin{pmatrix} \{+ + + +\} & \{+ + + -\} & \{- + + -\} & \{+ + - +\} \\ \{- - - +\} & \{- + + +\} & \{- - + +\} & \{- + - +\} \\ \{+ - - +\} & \{+ + - -\} & \{+ - - -\} & \{+ - + +\} \\ \{- - + -\} & \{+ - + -\} & \{- + - -\} & \{- - - -\} \end{pmatrix}. \tag{3}$$







$$\begin{pmatrix} -1 & -1 \\ -1 & -1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} -1 & -1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} -1 & -1 \\ -1 & -1 \end{pmatrix} \begin{pmatrix} -1 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \quad (14)$$

$$\begin{pmatrix} -1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} -1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ -1 & -1 \end{pmatrix} \begin{pmatrix} -1 & 1 \\ -1 & -1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -1 & -1 \end{pmatrix}. \quad (15)$$

of these, eight matrices are:

$$\begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ -1 & -1 \end{pmatrix} \begin{pmatrix} -1 & -1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} -1 & 1 \\ -1 & -1 \end{pmatrix} \begin{pmatrix} -1 & -1 \\ -1 & 1 \end{pmatrix} \quad (16)$$

are Hadamard matrices, because they satisfy the condition

$$H(2) \otimes H^T(2) = 2 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (17)$$

When raising any of the matrices (16) to Kronecker powers, the Hadamard matrices  $H(n)$  are again obtained, satisfying the condition:

$$H(n) \otimes H^T(n) = nI, \quad (18)$$

where  $I$  is an  $n \times n$  diagonal identity matrix:

$$I = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (19)$$

For example,

$$H(2)^{\otimes 2} = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}^{\otimes 2} = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} = \begin{pmatrix} 1 \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} & 1 \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \\ 1 \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} & -1 \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix} \quad (20)$$

$$H(2)^{\otimes 3} = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}^{\otimes 3} = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 & 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 \\ 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 \\ 1 & -1 & 1 & -1 & -1 & 1 & -1 & 1 \\ 1 & 1 & -1 & -1 & -1 & -1 & 1 & 1 \\ 1 & -1 & -1 & 1 & -1 & 1 & 1 & -1 \end{pmatrix} \quad (21)$$

and so on according to the algorithm

$$H(2)^{\otimes k} = H(2^k) = H(2) \otimes H(2)^{\otimes k-1} = H(2) \otimes H(2^{k-1}), \quad (22)$$

Hadamard matrices are used to construct error-correcting codes. In particular, Hadamard matrices are used to decipher the genetic code (Petoukhov, 2008; Petoukhov, 2020).

If in the matrices (20) and (21) again instead of 1 and  $-1$  we use the signs  $\{+\}$  and  $\{-\}$ , then we obtain the rule for raising two-row signatures to the Kronecker power, for example,

$$\begin{Bmatrix} + & + \\ + & - \end{Bmatrix}^{\otimes 2} = \begin{Bmatrix} + & + \\ + & - \end{Bmatrix} \otimes \begin{Bmatrix} + & + \\ + & - \end{Bmatrix} = \begin{Bmatrix} + \begin{Bmatrix} + & + \\ + & - \end{Bmatrix} & + \begin{Bmatrix} + & + \\ + & - \end{Bmatrix} \\ + \begin{Bmatrix} + & + \\ + & - \end{Bmatrix} & - \begin{Bmatrix} + & + \\ + & - \end{Bmatrix} \end{Bmatrix} = \begin{Bmatrix} + & + & + & + \\ + & - & + & - \\ + & + & - & - \\ + & - & - & + \end{Bmatrix} \quad (23)$$

$$\begin{pmatrix} + & + \\ + & - \end{pmatrix}^{\otimes 3} = \begin{pmatrix} + & + \\ + & - \end{pmatrix} \otimes \begin{pmatrix} + & + & + & + \\ + & - & + & - \\ + & + & - & - \\ + & - & - & + \end{pmatrix} = \begin{pmatrix} + & + & + & + & + & + & + & + \\ + & - & + & - & + & - & + & - \\ + & + & - & - & + & + & - & - \\ + & - & - & + & + & - & - & + \\ + & - & + & - & - & - & - & - \\ + & + & + & - & - & - & - & - \\ + & + & - & - & - & - & + & + \\ + & - & - & + & - & + & + & - \end{pmatrix} \quad (24)$$

double row stignatures,

$$\begin{pmatrix} - & + \\ + & + \end{pmatrix} \begin{pmatrix} + & + \\ - & + \end{pmatrix} \begin{pmatrix} + & - \\ + & + \end{pmatrix} \begin{pmatrix} + & + \\ + & - \end{pmatrix} \begin{pmatrix} + & - \\ - & - \end{pmatrix} \begin{pmatrix} - & - \\ + & - \end{pmatrix} \begin{pmatrix} - & + \\ - & - \end{pmatrix} \begin{pmatrix} - & + \\ - & + \end{pmatrix} \quad (25)$$

corresponding to matrices (15) will be called two-row Hadamard stignatures.

### 5.7. Noise-immune coding of genetic information

It was shown in (Petoukhov, 2008; Petoukhov, 2020) that Hadamard matrices (15) or two-row stignatures (25) can be used as the basis for studying noise-resistant genetic information. Deoxyribonucleic acid (DNA) molecules are built from four chemical elements (nucleotides):

$$\begin{array}{llll} \text{Adenine} & - & \text{A} & \text{or} & \text{I} \\ \text{Guanine} & - & \text{G} & \text{or} & \text{H} \\ \text{Thymine} & - & \text{T} & \text{or} & \text{V} \\ \text{Cytosine} & - & \text{C} & \text{or} & \text{H}' \end{array} \quad (26)$$

These four nucleotides correspond to two bits of information or four binary stignatures (9)

A	00	{- -}
G	01	{- +}
T	10	{+ -}
C	11	{+ +}

(27)

Nucleotides (27) form: doublets, triplets and other more complex combinations of informational polymer, i.e. DNA molecules (Petoukhov, 2008; Petoukhov, 2020):

$$\begin{pmatrix} A & G \\ C & T \end{pmatrix}^2 = \begin{pmatrix} A & G \\ C & T \end{pmatrix} \begin{pmatrix} A & G \\ C & T \end{pmatrix} = \begin{pmatrix} AA & AG \\ AH' & AV \\ CA & CG \\ CC & CT \end{pmatrix} \begin{pmatrix} GA & GG \\ GC & GT \\ TA & TG \\ TC & TT \end{pmatrix} \quad (28)$$

$$\begin{pmatrix} A & G \\ C & T \end{pmatrix}^3 = \begin{pmatrix} \begin{pmatrix} AAA & AAG \\ AAC & AAT \end{pmatrix} \begin{pmatrix} AGA & AGG \\ AGC & AGT \end{pmatrix} & \begin{pmatrix} GAA & GAG \\ GAC & GAT \end{pmatrix} \begin{pmatrix} GGA & GGG \\ GGC & GGT \end{pmatrix} \\ \begin{pmatrix} ACA & ACG \\ ACC & ACT \end{pmatrix} \begin{pmatrix} ATA & ATG \\ ATC & ATT \end{pmatrix} & \begin{pmatrix} GCA & GCG \\ GCC & GCT \end{pmatrix} \begin{pmatrix} GTA & GTG \\ GTC & GTT \end{pmatrix} \\ \begin{pmatrix} CAA & CAG \\ CAC & CAT \end{pmatrix} \begin{pmatrix} CGA & CGG \\ CGC & CGT \end{pmatrix} & \begin{pmatrix} TAA & TAG \\ TAC & TAT \end{pmatrix} \begin{pmatrix} TGA & TGG \\ TGC & TGT \end{pmatrix} \\ \begin{pmatrix} CCA & CCG \\ CCC & CCT \end{pmatrix} \begin{pmatrix} CTA & CTG \\ CTC & CTT \end{pmatrix} & \begin{pmatrix} TCA & TCG \\ TCC & TCT \end{pmatrix} \begin{pmatrix} TTA & TTG \\ TTC & TTT \end{pmatrix} \end{pmatrix} \quad (29)$$

These combinations of nucleotides correspond to the combinations of the "+" and "-" signs obtained from the two-row Hadamard signatures (25), for example

$$\left\{ \begin{matrix} + & + \\ + & - \end{matrix} \right\}^{\otimes 2} = \left\{ \begin{matrix} + & + & + & + \\ + & - & + & - \\ + & + & - & - \\ + & - & - & + \end{matrix} \right\}, \quad \left\{ \begin{matrix} + & + \\ + & - \end{matrix} \right\}^{\otimes 3} = \left\{ \begin{matrix} + & + & + & + & + & + & + & + \\ + & - & + & - & + & - & + & - \\ + & + & - & - & + & + & - & - \\ + & - & - & + & + & - & - & + \\ + & + & + & + & - & - & - & - \\ + & - & + & - & - & + & - & + \\ + & + & - & - & - & - & + & + \\ + & - & - & + & - & + & + & - \end{matrix} \right\}. \quad (30)$$

Thus, the Algebra of Signatures can be used to develop the theory of matrix coding of the space-time continuum.

### 5.8. Binary-coded decimal and Arithmetic of Signature

There is a complete correspondence between the binary-coded decimal (BCD) of digits and the 16 affine space signatures from the matrix (3):

Table 2: Relationship between digits, BCD and signatures

Digit	Binary-coded decimal				Signature
Allowed combinations					
	2 <sup>3</sup>	2 <sup>2</sup>	2 <sup>1</sup>	2 <sup>0</sup>	
0	0	0	0	0	{-----}
1	0	0	0	1	{----+}
2	0	0	1	0	{---+-}
3	0	0	1	1	{--++}
4	0	1	0	0	{-+--}
5	0	1	0	1	{-+-+}
6	0	1	1	0	{-++-}
7	0	1	1	1	{-+++}
8	1	0	0	0	{+---}
9	1	0	0	1	{+--}
Additional characters					
* (asterisk)	1	0	1	0	{+--+}
# (lattice)	1	0	1	1	{+ - + +}
+ (plus)	1	1	0	0	{+ + - -}
- (minus)	1	1	0	1	{+ + - +}
, (comma)	1	1	1	0	{+ + + -}
damping	1	1	1	1	{+ + + +}

From Table 2 it follows that if 0 is replaced by "-", and 1 is replaced by "+", then digits from matrix (3) can be assigned numbers. In addition, by analogy with binary-decimal arithmetic, the arithmetic of signatures can also be constructed, as shown in the example in Table 3.

Table 3: An example of an arithmetic operation for adding signatures

BCD addition operation example	Example of signature addition operation:
<p>Find the number <math>A = D + C</math>, where <math>D = 3927</math>, <math>C = 4856</math>.                      Let's represent the numbers <math>D</math> and <math>C</math> in binary decimal form:  <math>D = 3927_{10} = 0011\ 1001\ 0010\ 0111_{BCD}</math>  <math>C = 4856_{10} = 0100\ 1000\ 0101\ 0110_{BCD}</math>                      We sum the numbers <math>D</math> and <math>C</math> according to the rules of binary arithmetic:</p> $\begin{array}{r} \phantom{00}0011\ 1001\ 0010\ 0111 \\ + 0100\ 1000\ 0101\ 0110 \\ \hline = 1000\ 0001\ 0111\ 1101 - \text{Binary sum} \\ + \phantom{00}0110\phantom{00}0110 - \text{Correction} \\ \hline 1000\ 0111\ 1000\ 0011 \end{array}$ <p>where                      * – tetrad from which there was a transfer to the senior tetrad;                      ** – tetrad with a forbidden combination of bits.</p>	<p>Find the number <math>A = D + C</math>, where <math>D = 3927</math>, <math>C = 4856</math>.                      Let's represent the numbers <math>D</math> and <math>C</math> in signature form:  <math>D = 3927_{10} = \{- - + +\} \{+ - - +\} \{- - + -\} \{- + + +\}_{ASt}</math>  <math>C = 4856_{10} = \{- + - -\} \{+ - - -\} \{- + - +\} \{- + + -\}_{ASt}</math>                      We sum the numbers <math>D</math> and <math>C</math> according to the rules of signature arithmetic:</p> $\begin{array}{r} \phantom{00}\{- - + +\} \{+ - - +\} \{- - + -\} \{- + + +\} \\ + \{- + - -\} \{+ - - -\} \{- + - +\} \{- + + -\} \\ \hline = \{+ - - -\} \{- - - +\} \{- + + +\} \{+ + - +\} - \text{Binary sum} \\ \phantom{00}\{- + + -\} \phantom{00}\{- + + -\} \phantom{00}\{- + + -\} - \text{Correction} \\ \hline \{+ - - -\} \{- + + +\} \{+ - - -\} \{- - + +\} \end{array}$ <p>where                      * – tetrad from which there was a transfer to the senior tetrad;                      ** – tetrad with a forbidden combination of bits.</p>

Arithmetic of Signature can be used to describe the nodal configurations of affine spaces.

5.9. "Colored" quaternions

Sixteen signatures (3) correspond to 16 types of "colored" quaternions:

$z_1 = x_0 + ix_1 + jx_2 + kx_3$	$\{+ + + +\}$	$\{- - - -\}$	$z_9 = -x_0 - ix_1 - jx_2 - kx_3$	(31)
$z_2 = -x_0 - ix_1 - jx_2 + kx_3$	$\{- - - +\}$	$\{+ + + -\}$	$z_{10} = x_0 + ix_1 + jx_2 - kx_3$	
$z_3 = x_0 - ix_1 - jx_2 + kx_3$	$\{+ - - +\}$	$\{- + + -\}$	$z_{11} = -x_0 + ix_1 + jx_2 - kx_3$	
$z_4 = -x_0 - ix_1 + jx_2 - kx_3$	$\{- - + -\}$	$\{+ + - +\}$	$z_{12} = x_0 + ix_1 - jx_2 + kx_3$	
$z_5 = x_0 + ix_1 - jx_2 - kx_3$	$\{+ + - -\}$	$\{- - + +\}$	$z_{13} = -x_0 - ix_1 + jx_2 + kx_3$	
$z_6 = -x_0 + ix_1 - jx_2 - kx_3$	$\{- + - -\}$	$\{+ - + +\}$	$z_{14} = x_0 - ix_1 + jx_2 + kx_3$	
$z_7 = x_0 - ix_1 + jx_2 - kx_3$	$\{+ - + -\}$	$\{- + - +\}$	$z_{15} = -x_0 + ix_1 - jx_2 + kx_3$	
$z_8 = -x_0 + ix_1 + jx_2 + kx_3$	$\{- + + +\}$	$\{+ - - -\}$	$z_{16} = x_0 - ix_1 - jx_2 - kx_3$	

The complete set of "colored" quaternions (31) can be applied in various branches of mathematics and physics. By direct calculation it is easy to verify that the sum of all 16 types of "colored" quaternions (31) is equal to zero  $\sum_{k=1}^{16} z_k = 0$ . (32)

Expression (32) shows that the superposition (addition) of all types of "colored" quaternions is balanced with respect to zero, i.e. satisfies the "vacuum balance" condition.

5.10. Classes of signature

The signatures included in the matrix (3) can be divided into three classes:

Class 1: all signs in the signature are the same - 2 signatures:

$$\{+ + + +\} \{- - - -\}. \tag{33}$$

Class 2: two identical signs in a signature - 6 signatures:

$$\begin{aligned} &\{+ - - +\} \{+ + - -\} \{+ - + -\} \\ &\{- + + -\} \{- - + +\} \{- + - +\}. \end{aligned} \tag{34}$$

Class 3: three identical characters in a signature - 8 signatures:

$$\begin{aligned} &\{- - - +\} \{- - + -\} \{- + - -\} \{+ - - -\} \\ &\{+ + + -\} \{+ + - +\} \{+ - + +\} \{- + + +\}. \end{aligned} \tag{35}$$

5.11. Chess analogy

The chessboard has  $8 \times 8 = 64$  cells: 32 of them are black and 32 are white (see Fig. 9). Also, in the matrix of signatures (3) there are  $16 \times 4 = 64$  characters, of which 32 are plus "+" and 32 minus "-":

$$\begin{pmatrix} II & HI & VI & H'I \\ IH & HH & VH & H'H \\ IV & HV & VV & H'V \\ IH' & HH' & VH' & H'H' \end{pmatrix} \equiv \begin{pmatrix} \{++++\} & \{+++-\} & \{+-++\} & \{+--+ \} \\ \{++-+\} & \{++--\} & \{+--+\} & \{+---\} \\ \{-+++ \} & \{-+-+\} & \{-+ + +\} & \{- - + -\} \\ \{-+ - +\} & \{-+ - -\} & \{- - - +\} & \{- - - -\} \end{pmatrix}.$$




Fig. 9: The chessboard

At the beginning of the game, there are 32 chess pieces on the chessboard: 16 white and 16 black (see Fig. 9). Also, in the matrix (3) there are 16 signatures that coincide in properties with chess pieces in full accordance with their three classes (33) – (35) (see Fig. 10):

{+ - + +} pawn	{- - - +} pawn	{+ + - +} pawn	{+ - - -} pawn	{+ + + -} pawn	{- + + +} pawn	{- - + -} pawn	{- + - -} pawn
{- - + +} rook	{+ - + -} knight	{- + + -} bishop	{+ + + +} queen	{- - - -} king	{+ - - +} bishop	{- + - +} knight	{+ + - -} rook

Fig. 10: Illustration of the full correspondence of the 16 signatures of the matrix (3) to the 16 chess pieces

Matrix (3) can describe both "light" (i.e., light  $3D_{m,r}$  landscapes), which corresponds to 16 white chess pieces, and "darkness" (emptiness), which corresponds to 16 black chess pieces.

5.12. I Ching analogy

The I Ching (Chinese Book of Changes) is based on two Beginnings:

Yang            Yin  
 «—»    and    «-->»,

in the Algebra of Stignatures there are also two initial signs:

« - »    and    « + ».

The Book of Changes uses 8 trigrams (see Fig. 11a). Similarly, there are eight 3-bases in the Algebra of Stignatures (see Fig. 6a) with the following stignature

$$\begin{matrix} \{+ + +\} & \{- + +\} & \{- - +\} & \{- - -\} \\ \{+ - -\} & \{- + -\} & \{+ - +\} & \{+ + -\}, \end{matrix} \quad (36)$$

and eight 3-antibases (Fig. 6b) with opposite stignatures

$$\begin{matrix} \{- - -\} & \{+ - -\} & \{+ + -\} & \{+ + +\} \\ \{- + +\} & \{+ - +\} & \{- + -\} & \{- - +\}. \end{matrix} \quad (37)$$

In the Book of Changes, all possible combinations of two trigrams generate 64 hexagrams (Fig. 11b), and in the Algebra of Stignatures, 64 combinations of each 3-basis with stignatures (36) with each 3-antibasis with opposite stignatures (37) are possible.

The three-logic of the Book of Changes is consistent with the four-logic of the Stignature Algebra if in the ranking expression (5) one column of signs is transferred to another side of equalities:

$$\begin{matrix} 0 = & \{0 & 0 & 0\} & + & \{0 & 0 & 0\} & = & 0 \\ - = & \{+ & + & +\} & + & \{- & - & -\} & = & + \\ + = & \{- & - & +\} & + & \{+ & + & -\} & = & - \\ - = & \{- & - & +\} & + & \{+ & + & -\} & = & + \\ + = & \{- & + & -\} & + & \{+ & - & +\} & = & - \\ - = & \{+ & - & -\} & + & \{- & + & +\} & = & + \\ + = & \{+ & - & -\} & + & \{- & + & +\} & = & - \\ - = & \{- & + & -\} & + & \{+ & - & +\} & = & + \\ + = & \{+ & + & +\} & + & \{- & - & -\} & = & - \\ 0 = & \{0 & 0 & 0\}_+ & + & \{0 & 0 & 0\}_+ & = & 0 \end{matrix} \quad (38)$$

The Book of Changes also uses four combinations of two principles:

«—»    and    «-->»:

Young Yin	Old Yin	Young Yang	Old Yang
--	—	--	—
—	--	--	—
<i>Binary stignatures</i>			
{+ -}	{- +}	{--}	{+ +}





Fig. 11: Trigrams and hexagrams of the I Ching (Book of Changes)

Similarly, in the Algebra of Signatures, four binary combinations of signs "+" and "-" (9) are possible: {+ +}, {- -}, {+ -} {- +}, from which affine space signatures (11) are formed, etc.

### 5.13. Hebrew analogy

Judaism is based on Algorithms for revealing the Great Name of the GOD יה-וה-י (Gaukhman, 2007a; Gaukhman, 2007b). Further, instead of Hebrew letters, the transliteration יה-וה-י ≡ H' V H I will be used.

One of the endless Revelations of the Name H' V H I is called "Etz Chaim" ("Tree of Life" or "Tree of the Sefirot"):

$$\begin{pmatrix} I & H \\ H' & V \end{pmatrix}^{\otimes 2} = \begin{pmatrix} I \begin{pmatrix} I & H \\ H' & V \end{pmatrix} & H \begin{pmatrix} I & H \\ H' & V \end{pmatrix} \\ H' \begin{pmatrix} I & H \\ H' & V \end{pmatrix} & V \begin{pmatrix} I & H \\ H' & V \end{pmatrix} \end{pmatrix} = \begin{pmatrix} \begin{pmatrix} II & IH \\ IH' & IV \end{pmatrix} & \begin{pmatrix} HI & HH \\ HH' & HV \end{pmatrix} \\ \begin{pmatrix} H'I & H'H \\ H'H' & H'V \end{pmatrix} & \begin{pmatrix} VI & VH \\ VH' & VV \end{pmatrix} \end{pmatrix}. \tag{39}$$

The components of this matrix correspond to 10 Sefirot (i.e. Qualities) of Spirituality:

Letter of the Name יה-וה-י	Component of the Matrix of Qualities (39) (see Fig. 12)	Sefirah
<i>i</i> tip of the Letter Yud	II	Keter
I	HH	Hochma
H	VV	Bina
V	IV, IH, IH', VH, VH', HH', VI, HI, H'I, HV, H'V, H'H	Tiphereth*
H'	H'H'	Malchut




Fig. 12: Tree of ten Sefirot (Qualities)

where the Sefira Tipheret\* consists of six dual Sefirot [13]:

Chesed ( $IV = -VI$ ), Gevura ( $IH = -HI$ ), Tiferet ( $IH' = -H'I$ ), Netzach ( $VH = -HV$ ), Hod ( $VH' = -V'H$ ), Yesod ( $HH' = -H'H$ ).

The Qualities Matrix (39) can be written as the sum of two matrices (Gaukhman, 2007a; Gaukhman, 2007b).

$$\begin{pmatrix} II & 0 & 0 & 0 \\ 0 & HH & 0 & 0 \\ 0 & 0 & VV & 0 \\ 0 & 0 & 0 & H'H' \end{pmatrix} + \begin{pmatrix} 0 & HI & VI & H'I \\ IH & 0 & VH & H'H \\ IV & HV & 0 & H'V \\ IH' & HH' & VH' & 0 \end{pmatrix}. \tag{40}$$

Similarly, the signature matrix (3) is the result of raising the two-row matrix to the second Kronecker power



$$\begin{pmatrix} I & H \\ H' & V \end{pmatrix}^{\otimes 2} = \begin{pmatrix} \{++\} & \{+-\} \\ \{-+\} & \{--\} \end{pmatrix}^{\otimes 2} = \begin{pmatrix} \{++++\} & \{+++-\} & \{+-++\} & \{+--+ \} \\ \{++-+\} & \{++--\} & \{+---\} & \{+----\} \\ \{-+++ \} & \{-++-\} & \{---+\} & \{----\} \\ \{-+-+\} & \{-+--\} & \{----\} & \{-----\} \end{pmatrix}.$$

This matrix can also be represented as the sum of the diagonal and antisymmetric matrices (13)

$$\begin{pmatrix} \{++++\} & 0 & 0 & 0 \\ 0 & \{-+++ \} & 0 & 0 \\ 0 & 0 & \{+---\} & 0 \\ 0 & 0 & 0 & \{-----\} \end{pmatrix} + \begin{pmatrix} 0 & \{+++-\} & \{-++-\} & \{++-+\} \\ \{----+\} & 0 & \{---+\} & \{-+-+\} \\ \{+---+\} & \{+---\} & 0 & \{+-++\} \\ \{-+-+\} & \{+-+--\} & \{-+--\} & 0 \end{pmatrix}. \tag{41}$$

Comparing the matrices (40) and (41), we find that the signatures of the matrix (3) can reflect the characteristic properties of the corresponding Sefirot (Qualities) of the "Tree of Life" (Gaukhman, 2007a; Gaukhman, 2007b).

$$\begin{pmatrix} II & 0 & 0 & 0 \\ 0 & HH & 0 & 0 \\ 0 & 0 & VV & 0 \\ 0' & 0 & 0 & H'H' \end{pmatrix} \equiv \begin{pmatrix} \{++++\} & 0 & 0 & 0 \\ 0 & \{-+++ \} & 0 & 0 \\ 0 & 0 & \{+---\} & 0 \\ 0 & 0 & 0 & \{-----\} \end{pmatrix}, \tag{42}$$

$$\begin{pmatrix} 0 & HI & VI & H'I \\ IH & 0 & VH & H'H \\ IV & HV & 0 & H'V \\ IH' & HH' & VH' & 0 \end{pmatrix} \equiv \begin{pmatrix} 0 & \{+++-\} & \{-++-\} & \{++-+\} \\ \{----+\} & 0 & \{---+\} & \{-+-+\} \\ \{+---+\} & \{+---\} & 0 & \{+-++\} \\ \{-+-+\} & \{+-+--\} & \{-+--\} & 0 \end{pmatrix}.$$

Thus, the following analogy can be traced between the ten Sephira (Quality) and the sixteen signatures of the matrix (3):

- 1)  $\{++++\} + \{+---\} = 0$  - expression of the properties of Sefira Keter II and Sefira Malchut H'H';
- 2)  $\{-+++ \} + \{+---\} = 0$  - expression of the properties of Sefira Hochma VV and Sefira Binah HH;
- 3)  $\{---+\} + \{++-+\} = 0$  - expression of the properties of the Sefira Gevura (IH = - HI);
- 4)  $\{+--+\} + \{-+-+\} = 0$  - expression of the properties of the Sefira Chesed (IV = - VI);
- 5)  $\{---+\} + \{++-+\} = 0$  - expression of the properties of the Sefira Teferet (IH' = - H'I);
- 6)  $\{-+-+\} + \{+-++\} = 0$  - expression of properties of Sefira Netzach (VH' = - H'V);
- 7)  $\{+-+--\} + \{-+-+\} = 0$  - expression of the properties of the Sephirah Yesod (HH' = - H'H);
- 8)  $\{+-+--\} + \{-+-+\} = 0$  - expression of the properties of the Sefira Hod (HV = - VH).

This analogy corresponds to the criteria of "Zero Philosophy", emanating from "EIN SOF, Baruhu" (INFINITE NOTHING, Blessed be HE). In turn, from the "Zero Philosophy" follows the condition of "vacuum balance" and the structure of "split zero" (5), on which the geometrized vacuum physics and, in particular, the Algebra of Stignatures are based.

Let's return to the representation of the Great Name of GOD יה-וה-י in the form of binary stignatures (9)

$$\text{יה-וה-י} \equiv H' V H I \equiv H' \leftrightarrow \{+ -\}, V \leftrightarrow \{- +\}, H \leftrightarrow \{++\}, I \leftrightarrow \{--\}. \tag{42b}$$



Let's transfer the stignatures  $\{- + + +\}$  and  $\{+ - - -\}$  from the numerators of the ranks (43) to their denominators with inversion, i.e. with signs reversed. Such an operation does not violate the "vacuum balance" and leads to the ranking expression

$$\begin{array}{lcl}
 \{+ + + +\} & + & \{- - - -\} = 0 \\
 \{- - - +\} & + & \{+ + + -\} = 0 \\
 \{+ - - +\} & + & \{- + + -\} = 0 \\
 \{- - + -\} & + & \{+ + - +\} = 0 \\
 \{+ + - -\} & + & \{- - + +\} = 0 \\
 \{- + - -\} & + & \{+ - + +\} = 0 \\
 \underline{\{+ - + -\}} & + & \underline{\{- + - +\}} = 0 \\
 \{+ - - -\}_+ & + & \{- + + +\}_+ = 0 .
 \end{array} \tag{44}$$

In this case, in the denominator of the left rank in Ex. (44), the stignature of the affine 4-Minkowski space  $\{+ - - -\}$  was obtained, and in the denominator of the right rank in Ex. (44), the stignature of the affine 4-anti-Minkowski space  $\{- + + +\}$  (Gaukhman, 2007b) was obtained.

Not only the ranking expression (44) leads to a balanced (with respect to zero) dyad of affine Minkowski stignatures

$$\{+ - - -\} + \{- + + +\} = 0.$$

The following ranking expressions that preserve the vacuum balance lead to the same result:

$$\begin{array}{lcl}
 \{- - - +\} & + & \{+ + + -\} = 0 \\
 \{+ - + -\} & + & \{- + - +\} = 0 \\
 \underline{\{+ + - -\}} & + & \underline{\{- - + +\}} = 0 \\
 \{+ - - -\}_+ & + & \{- + + +\}_+ = 0
 \end{array} \tag{45}$$

$$\begin{array}{lcl}
 \{- - + -\} & + & \{+ + - +\} = 0 \\
 \{+ + - -\} & + & \{- - + +\} = 0 \\
 \underline{\{+ - - +\}} & + & \underline{\{- + + -\}} = 0 \\
 \{+ - - -\}_+ & + & \{- + + +\}_+ = 0
 \end{array} \tag{46}$$

$$\begin{array}{lcl}
 \{- - + -\} & + & \{+ + - +\} = 0 \\
 \{+ + - -\} & + & \{- - + +\} = 0 \\
 \underline{\{+ - - +\}} & + & \underline{\{- + + -\}} = 0 \\
 \{+ - - -\}_+ & + & \{- + + +\}_+ = 0
 \end{array} \tag{47}$$

The Algebra of Stignature can be used as a musical score of spatial harmony. Just like in music, stignatures form balanced octaves ( $7+1=8$ ) of the form (44) and chords ( $3+1=4$ ) of the form (45) – (47). Moreover, in different situations, the Algebra of Stignatures uses a different number of signs "+" and "-" in proportions in relation to the total number: 1, 1/2, 1/8, 1/16, 1/32, 1/64, etc. This corresponds to the proportions of the duration of the notes: 1, 1/2, 1/8, 1/16, 1/32, 1/64, etc.

Ancient philosophers, beginning with the Pythagoreans, believed that Space was filled with harmonious musical rhythms and proportions. Music theory was later used by Johannes Kepler in «*Harmonices Mundi*» (Harmony of the World), which led to the discovery of the third law of celestial mechanics. It is possible that the combination of musical techniques with the Algebra of Stignatures algorithms will also lead to interesting results.

### 5.15. Pythagorean analogy (Algebra of Signatures)

Let's pass from affine spaces to metric spaces. For this, as an example, we use an affine space with a 4-basis  $\mathbf{e}_i^{(7)}(\mathbf{e}_0^{(7)}, \mathbf{e}_1^{(7)}, \mathbf{e}_2^{(7)}, \mathbf{e}_3^{(7)})$  (see Figs. 7 and 13) with signature  $\{+ + + -\}$ . In this space, we define a 4-vector

$$d\mathbf{s}^{(7)} = \mathbf{e}_i^{(7)} dx^{i(7)} = \mathbf{e}_0^{(7)} dx^{0(7)} + \mathbf{e}_1^{(7)} dx^{1(7)} + \mathbf{e}_2^{(7)} dx^{2(7)} + \mathbf{e}_3^{(7)} dx^{3(7)}, \quad (48)$$

where  $dx_i^{(7)}$  is the  $i$ -th projection of the 4-vector  $d\mathbf{s}^{(7)}$  onto the  $x_i^{(7)}$  axis, the direction of which is determined by the basis vector  $\mathbf{e}_i^{(7)}$ .




Fig. 13: Two 4-bases with different stignatures

Similarly, we define the second 4-vector in the affine space with the 4-basis  $\mathbf{e}_i^{(5)}(\mathbf{e}_0^{(5)}, \mathbf{e}_1^{(5)}, \mathbf{e}_2^{(5)}, \mathbf{e}_3^{(5)})$  (Figs. 7 and 13), with signature  $\{+ + + +\}$

$$d\mathbf{s}^{(5)} = \mathbf{e}_i^{(5)} dx^{i(5)} = \mathbf{e}_0^{(5)} dx^{0(5)} + \mathbf{e}_1^{(5)} dx^{1(5)} + \mathbf{e}_2^{(5)} dx^{2(5)} + \mathbf{e}_3^{(5)} dx^{3(5)}. \quad (49)$$

We find the scalar product of 4-vectors (48) and (49)

$$\begin{aligned} ds^{(5,7)2} &= d\mathbf{s}^{(5)} d\mathbf{s}^{(7)} = \mathbf{e}_i^{(5)} \mathbf{e}_j^{(7)} dx^i dx^j = \\ &= \mathbf{e}_0^{(5)} \mathbf{e}_0^{(7)} dx^0 dx^0 + \mathbf{e}_1^{(5)} \mathbf{e}_0^{(7)} dx^1 dx^0 + \mathbf{e}_2^{(5)} \mathbf{e}_0^{(7)} dx^2 dx^0 + \mathbf{e}_3^{(5)} \mathbf{e}_0^{(7)} dx^3 dx^0 + \\ &+ \mathbf{e}_0^{(5)} \mathbf{e}_1^{(7)} dx^0 dx^1 + \mathbf{e}_1^{(5)} \mathbf{e}_1^{(7)} dx^1 dx^1 + \mathbf{e}_2^{(5)} \mathbf{e}_1^{(7)} dx^2 dx^1 + \mathbf{e}_3^{(5)} \mathbf{e}_1^{(7)} dx^3 dx^1 + \\ &+ \mathbf{e}_0^{(5)} \mathbf{e}_2^{(7)} dx^0 dx^2 + \mathbf{e}_1^{(5)} \mathbf{e}_2^{(7)} dx^1 dx^2 + \mathbf{e}_2^{(5)} \mathbf{e}_2^{(7)} dx^2 dx^2 + \mathbf{e}_3^{(5)} \mathbf{e}_2^{(7)} dx^3 dx^2 + \\ &+ \mathbf{e}_0^{(5)} \mathbf{e}_3^{(7)} dx^0 dx^3 + \mathbf{e}_1^{(5)} \mathbf{e}_3^{(7)} dx^1 dx^3 + \mathbf{e}_2^{(5)} \mathbf{e}_3^{(7)} dx^2 dx^3 + \mathbf{e}_3^{(5)} \mathbf{e}_3^{(7)} dx^3 dx^3. \end{aligned} \quad (50)$$

For the case under consideration, the scalar products of basis vectors  $\mathbf{e}_i^{(5)} \mathbf{e}_j^{(7)}$  are:

$$\begin{aligned} \text{for } i = j \quad \mathbf{e}_0^{(5)} \mathbf{e}_0^{(7)} = 1, \quad \mathbf{e}_1^{(5)} \mathbf{e}_1^{(7)} = 1, \quad \mathbf{e}_2^{(5)} \mathbf{e}_2^{(7)} = 1, \quad \mathbf{e}_3^{(5)} \mathbf{e}_3^{(7)} = -1, \\ \text{for } i \neq j, \quad \text{all } \mathbf{e}_i^{(5)} \mathbf{e}_j^{(7)} = 0. \end{aligned} \quad (51)$$

In this case, expression (50) takes the form of a quadratic form

$$ds^{(5,7)2} = dx^0 dx^0 + dx^1 dx^1 + dx^2 dx^2 - dx^3 dx^3 = dx^{02} + dx^{12} + dx^{22} - dx^{32} \quad (52)$$

with signature  $(+ + + -)$ .

"Signature" (GRT term) is an ordered set of signs in front of the corresponding terms of the quadratic form.

To determine the signature of the metric space with the metric (52), instead of performing the scalar product of vectors (50), it is enough to multiply the signs of the 4-basis signatures shown in Fig. 18 by columns:

$$\frac{\begin{matrix} \{+ + + +\} \\ \{+ + + -\} \\ (+ + + -)_\times \end{matrix}}{\hspace{10em}} \quad (53)$$

In the numerator of the rank (53), the signs in each column are multiplied according to the rules

$$\{+\} \times \{+\} = \{+\}; \quad \{-\} \times \{+\} = \{-\}; \quad (54)$$

the result of such multiplication is written in the denominator (under the line) of the same column. The execution of actions according to these rules will be called rank multiplication.

Just as it was done with the vectors  $ds^{(5)}$  and  $ds^{(7)}$  {see expressions (48) – (53)}, pairwise scalarly multiply vectors from all 16 affine spaces with 4 bases shown in Fig. 7. As a result, we obtain  $16 \times 16 = 256$  metric 4-spaces with 4-intervals of the form

$$ds^{(ab)2} = e_i^{(a)} e_j^{(b)} dx^{i(a)} dx^{j(b)}, \quad (55)$$

where  $a = 1, 2, 3, \dots, 16$ ;  $b = 1, 2, 3, \dots, 16$ .

The signatures of these  $16 \times 16 = 256$  metric 4-spaces can be defined, similarly to (53), by rank multiplications of the signs of the signatures corresponding to the affine spaces, for example:

$$\frac{\begin{matrix} \{+ - + +\} & \{+ + + +\} & \{- + + +\} & \{+ + + +\} \\ \{+ + + -\} & \{+ - + -\} & \{+ + + -\} & \{- + + -\} & \dots \\ (+ - + -)_\times & (+ - + -)_\times & (- + + -)_\times & (- + + -)_\times \end{matrix}}{\hspace{10em}}$$

Thus, point  $O$  (see Fig. 5) is the place of intersection of all 256 metric 4-spaces with metrics (55) and signatures (40). In this variety of intersecting metric spaces, there are many features and patterns that are studied by the Algebra of Signatures.

The fundamentals of the Algebra of Signature will be covered in the next article of this project. Here we only note that quadratic forms (i.e., metrics) of type (52), or

$$ds^2 = dx_0^2 + dx_1^2 + dx_2^2 + dx_3^2 \quad (57)$$

originate from the Pythagorean theorem  $c^2 = a^2 + b^2$ , the proof of which in about 520 BC, brought about the greatest revolution in the consciousness of mankind.

### 6 Spectra-stignature analysis

As an example, we use the Algebra of Signatures to expand the possibilities of spectral analysis.

Let's recall the well-known in quantum physics procedure for the transition from the coordinate representation to the momentum one. Let the probability distribution density of the location of an elementary particle be given as a function of space and time  $\rho(ct, x, y, z)$ . This function is represented as a product of two amplitudes:

$$\rho(ct, x, y, z) = \varphi(ct, x, y, z) \varphi(ct, x, y, z). \quad (58)$$

Next, two Fourier transforms are performed

$$\psi(p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi(ct, x, y, z) \exp\{i \frac{p}{\eta} (ct - x - y - z)\} d\Omega \quad \text{signature } \{+ - - -\}, \quad (59)$$

$$\psi^*(p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi(ct, x, y, z) \exp\{i \frac{p}{\eta} (-ct + x + y + z)\} d\Omega \quad \text{signature } \{- + + +\}, \quad (60)$$

where  $p = 2\pi\eta/\lambda$  is the generalized frequency;  $\lambda$  is the wavelength;  $\eta$  is the coefficient of proportionality (in quantum mechanics,  $\eta = \hbar$  is the reduced Planck constant);  $d\Omega = dctdx dy dz$  is an elementary 4-dimensional volume.

The momentum (i.e., spectral) representation of the function  $\rho(ct, x, y, z)$  is obtained as a result of the product of two probability amplitudes (59) and (60)

$$G(p_{ct}, p_x, p_y, p_z) = \psi(p_{ct}, p_x, p_y, p_z) \cdot \psi^*(p_{ct}, p_x, p_y, p_z). \quad (61)$$

By analogy with the procedure (59) - (60), we formulate the basics of the spectral signature analysis. We represent the function  $\rho(ct, x, y, z)$  as a product of not two, but 8 "amplitudes"

$$\rho(ct, x, y, z) = \varphi_1(ct, x, y, z) \varphi_2(ct, x, y, z) \varphi_3(ct, x, y, z) \times \dots \times \varphi_8(ct, x, y, z) = \prod_{k=1}^8 \phi_k(ct, x, y, z).$$

Instead of the imaginary unit  $i$ , which was used in the integrals (59) and (60), we introduce into consideration eight objects  $\zeta_r$  (where  $r = 1, 2, 3, \dots, 8$ ) satisfying the anticommutative relations of the Clifford algebra:

$$\zeta_m \zeta_k + \zeta_k \zeta_m = 0 \quad \text{for } m \neq k, \quad \zeta_m \zeta_m = 1, \quad (63)$$

$$\text{or } \zeta_m \zeta_k + \zeta_k \zeta_m = 2\delta_{km}, \quad (64)$$

where  $\delta_{km}$  is the Kronecker symbol ( $\delta_{km} = 0$  for  $m \neq k$  and  $\delta_{km} = 1$  for  $m = k$ ).

These requirements are met, for example, by a set of  $8 \times 8$  matrices of the type

$$\begin{aligned}
 \zeta_1 &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} & \zeta_2 &= \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \end{pmatrix} & \zeta_3 &= \begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix} \\
\zeta_4 &= \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix} & \zeta_5 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \end{pmatrix} & \zeta_6 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \\
\zeta_7 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} & \zeta_8 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} & \delta_{km} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}
 \end{aligned} \tag{65}$$

Let's perform eight "color" Fourier transforms:

	Stignature
$\psi_1(p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_1(ct, x, y, z) \exp\{\zeta_1 \frac{p}{\eta} (ct + x + y + z)\} d\Omega,$	{+ + + +}
$\psi_2(p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_2(ct, x, y, z) \exp\{\zeta_2 \frac{p}{\eta} (-ct - x - y + z)\} d\Omega,$	{- - - +}
$\psi_3(p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_3(ct, x, y, z) \exp\{\zeta_3 \frac{p}{\eta} (ct - x - y + z)\} d\Omega,$	{+ - - +}
$\psi_4(p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_4(ct, x, y, z) \exp\{\zeta_4 \frac{p}{\eta} (-ct - x + y - z)\} d\Omega,$	{- - + -}
$\psi_5(p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_5(ct, x, y, z) \exp\{\zeta_5 \frac{p}{\eta} (ct + x - y - z)\} d\Omega,$	{+ + - -}
$\psi_6(p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_6(ct, x, y, z) \exp\{\zeta_6 \frac{p}{\eta} (-ct + x - y - z)\} d\Omega,$	{- + - -}
$\psi_7(p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_7(ct, x, y, z) \exp\{\zeta_7 \frac{p}{\eta} (ct - x + y - z)\} d\Omega,$	{+ - + -}
$\psi_8(p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_8(ct, x, y, z) \exp\{\zeta_8 \frac{p}{\eta} (-ct + x + y + z)\} d\Omega,$	{- + + +}
	{0 0 0 0}

where the objects  $\zeta_m$ (60) perform the function of Clifford imaginary units.

We also find eight complex-conjugate Fourier images with opposite stignatures:

	Stignature
$\psi_1 * (p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_1(ct, x, y, z) \exp\{\zeta_1 \frac{p}{\eta} (-ct - x - y - z)\} d\Omega,$	{-- --}
$\psi_2 * (p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_2(ct, x, y, z) \exp\{\zeta_2 \frac{p}{\eta} (c + x + y - z)\} d\Omega,$	{+ + + -}
$\psi_3 * (p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_3(ct, x, y, z) \exp\{\zeta_3 \frac{p}{\eta} (-ct + x + y - z)\} d\Omega,$	{- + + -}
$\psi_4 * (p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_4(ct, x, y, z) \exp\{\zeta_4 \frac{p}{\eta} (ct + x - y + z)\} d\Omega,$	{+ + - +}
$\psi_5 * (p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_5(ct, x, y, z) \exp\{\zeta_5 \frac{p}{\eta} (-ct - x + y + z)\} d\Omega,$	{- - + +}
$\psi_6 * (p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_6(ct, x, y, z) \exp\{\zeta_6 \frac{p}{\eta} (ct - x + y + z)\} d\Omega,$	{+ - + +}
$\psi_7 * (p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_7(ct, x, y, z) \exp\{\zeta_7 \frac{p}{\eta} (-ct + x - y + z)\} d\Omega,$	{- + - +}
$\psi_8 * (p_{ct}, p_x, p_y, p_z) = \int_{-\infty}^{\infty} \phi_8(ct, x, y, z) \exp\{\zeta_8 \frac{p}{\eta} (ct - x - y - z)\} d\Omega.$	{+ - - -}
	{0 0 0 0}

(67)


The integrals of the "colored" Fourier transform (61) and (62) include 16 linear forms with matrix signatures (3). The spectral-stignature representation of the function  $\rho(ct, x, y, z)$  is obtained as a result of the product of eight corresponding pairs of "color" amplitudes (66) and their complex conjugate "color" amplitudes (67)

$$\Re(p_{ct}, p_x, p_y, p_z) = \prod_{k=1}^8 \psi_k(p_{ct}, p_x, p_y, p_z) \psi_k^* (p_{ct}, p_x, p_y, p_z). \tag{68}$$

The spectral-stignature Fourier analysis proposed here can be useful for the development of "zero" technologies, such as sealing vacuum communication channels, and many other branches of science and technology.

### Carthusian crucifix

The orthogonal Cartesian coordinate system is mentally connected with the crucifixion of Jesus Christ. Modern science was created by the universities of medieval Europe, which looked at the world around them through a crosshair with a suffering Savior. Also, the signs of stignatures from matrix (3) characterize the directions of the axes of the multidimensional crucifixion and are associated with the Stigmata (Greek *στιγματος* - signs, marks, ulcers, wounds) on the head, hands and feet of the crucified Messiah.



## CONCLUSIONS

This work is the first of a series of articles under the general title "Geometrized vacuum physics from the standpoint of the Algebra of Signatures", where, as a result, based on the study of vacuum properties, it is assumed:

- build a multilayer and multilevel cosmological model with the development of ideas about the nature of all known force interactions, including gravity;
- present metric-dynamic models of all elementary particles and quarks that are part of the Standard Model (with the exception of the Higgs boson);
- develop a theoretical foundation for the development of "zero" (i.e., vacuum) technologies, including vacuum ethics, aesthetics and energy.

This article proposes to explore a local area of ideal vacuum (or "vacuum", i.e. void) by probing it with light rays with a monochromatic wavelength  $\lambda_{m,n}$  from three mutually perpendicular directions. As a result, a 3-dimensional light cubic lattice is formed in the "vacuum", which is called the  $\lambda_{m,n}$ -vacuum (or 3D $_{m,n}$ -landscape) with the edge length of the cubic cell  $\varepsilon_{m,n} \sim 100 \cdot \lambda_{m,n}$  (see Fig. 1 and 2). If we similarly probe the same volume of vacuum with monochromatic light rays with other wavelengths  $\lambda_{m+k,n+k}$ , then we get an infinite number of  $\lambda_{m,n}$ -vacuums nested



into each other like nesting dolls (see Fig. 2). Such an infinite hierarchical discrete sequence of light  $3D_{m,r}$ -landscapes is called a longitudinal bundle of a 3-dimensional volume of "vacuum" (emptiness).

The study of the geometric features of one ideal cubic cell of any of the  $\lambda_{m,r}$ -vacuums led to the development of the Algebra of Stignature. An analysis of the properties of the matrix of stignatures (3) showed that the Algebra of Stignatures is a unique mathematical and logical apparatus that has its mental roots in ancient philosophical traditions.

The foundations of the Algebra of Stignature presented in this article can be applied not only in vacuum physics (as will be shown in the following articles of this project), but also in many other branches of knowledge, for example, in multidimensional geometry, topology, continuum mechanics, crystallography, spatial coding theory, algebraic genetics, relativity theory, etc. In this article, as an example, the use of the Algebra of Stignatures for the development of stignature-spectral analysis is presented.

## ACKNOWLEDGMENTS

I express my sincere gratitude to Gavriil Davydov, David Reid and Eliezer Rayhman for their assistance. The discussion of the article was attended by Academician of the Russian Academy of Sciences Shipov G.I., Ph.D. Lukyanov V.A., Lebedev V.A., Prokhorov S.G. and Khramikhin V.P. Also, the author is grateful for the support of Salova M.N., Morozova T.S., Przhigodsky S.V., Maslov A.N., Bolotov A.Yu., Ph.D. Levi T.S., Musanov S.V., Batanova L.A., Ph.D. Myshelov E.P., Chivikov E.P.

## REFERENCES

Barrow, J.D. (2002). *"The Book of Nothing: Vacuums, Voids, and the Latest Ideas About the Origins of the Universe"*. Vintage Series. Vintage. pp. 71–72, 77. ISBN 978-0-375-72609-5. LCCN 00058894.

Chambers, A. (2004). *"Modern Vacuum Physics"*. Boca Raton: CRC Press, 2004. — ISBN 0-8493-2438-6.

Di Teodoro, A. & Contreras, E. (2023). A vacuum solution of modified Einstein equations based on fractional calculus. *Eur. Phys. J. C* 83: 434, arXiv:2305.15232 Related DOI:<https://doi.org/10.1140/epjc/s10052-023-11626-4>

Gaukhman, M.Kh. (2007a). *Algebra of signatures "NAMES" (orange Alsigna)*. - M.: LKI, 2007, p.228, ISBN 978-5-382-00077-0, (available at [на www.alsigna.ru](http://www.alsigna.ru)) [in Russian].

Gaukhman, M.Kh. (2007b). *Algebra of signatures "Emptiness" (yellow Alsigna)*. - M.: URSS, 2007. - 308 pages. ISBN 978-5-382-00580-5, (available at [на www.alsigna.ru](http://www.alsigna.ru)).

Peatross J. & Ware M. (2015). *"Physics of Light and Optics"*. Brigham Young University, p. 338, ISBN 978-1-312-92927-2.

Petoukhov S.V. (2008). *"Matrix genetics, algebras of the genetic code, noise-immunity"*. Moscow: RCD, 316 p. ISBN 978-5-93972-643-6 <http://petoukhov.com/matrix-genetics-petoukhov-2008.pdf> (in Russian).

Petoukhov S.V. (2020). *"The rules of long DNA-sequences and tetra-groups of oligonucleotides"*. arXiv:1709.04943v6.

Roberts, M.D. (2000). *"Vacuum Energy"*. High Energy Physics". Theory: hep-th/0012062. arXiv:hep-th/0012062. Bibcode:2000hep.th...12062R.

Shipov, G. (1998). "*A Theory of Physical Vacuum*". Moscow ST-Center, Russia ISBN 5 7273-0011-8.

Tyurin, Yu.I., Chernov I.P. & Kryuchkov Yu.Yu. (2009). "*Optics*". Tomsk Polytechnic University Publishing House, p. 240, ISBN 5-98298-434-5 [in Russian].